

**Research Article****Evaluation of Antioxidant activity of *Cardiospermum halicacabum* Linn leaves in Atherodiet induced hyperlipidemic wistar albino rats****<sup>1\*</sup>Thamizh Selvam N, <sup>2</sup>Surabhi KR, <sup>3</sup>Sanjaya Kumar YR, <sup>4</sup>Vasanth Kumar KG, <sup>5</sup>Sudesh Gaidhani N, <sup>6</sup>Radhakrishnan P**<sup>1</sup>Assistant Director-Biochemistry, Department of Biochemistry and Pathology, National Ayurveda Research Institute for Panchakarma, Cheruthuruthy.<sup>2</sup>Senior Research Fellow, Department of Biochemistry and Pathology, NARIP, Cheruthuruthy<sup>3</sup>Assistant Director- Pharmacology, Department of Pharmacology, NARIP, Cheruthuruthy<sup>4</sup>Assistant Director- Chemistry, Department of Chemistry, NARIP Cheruthuruthy<sup>5</sup>Assistant Director-Pharmacology, Programme Officer, Central Council for Research in Ayurvedic Sciences, New Delhi<sup>6</sup>Assistant Director In-Charge, Head of the Institution, NARIP, Cheruthuruthy.

Received: 20 March 2018

Revised: 25 April 2018

Accepted: 3 May 2018

**Abstract**

**Objective:** *Cardiospermum halicacabum* Linn commonly known as 'Balloon vine' is a climber found in Western ghat region. The plant is claimed for medicinal properties and used by traditional healers of Kerala for treatment of various ailments. The present study has evaluated the antioxidant potential of *C. halicacabum* Linn Leaf in the atherodiet induced hyperlipidemic wistar albino rats. **Materials and methods:** The study comprised of six groups such as control group, test extract group, disease control group, treatment group low dose, treatment group high dose and standard drug group. The duration of the study was 28 days and the extract was administered in the animals as per the standard protocol. The antioxidant parameters such as Superoxide dismutase, Catalase, Glutathione and Glutathione peroxidase were evaluated in the blood and tissue samples of liver, kidney and heart. **Results and conclusion:** The study evidenced that the *C. halicacabum* leaf has significant antioxidant potential in blood and tissue level when compared with different groups.

**Keywords:** *C. halicacabum*, antioxidant activity, hyperlipidemia, SOD, catalase**Introduction**

The role of free radicals in causing of various diseases such as cancer, atherosclerosis, diabetes and other life style diseases have been well reported by scientific communities (Pawan et al., 2011; Harman, 1992). So, the studies related on scavenging of free radicals through antioxidants have become one of the core research areas in health science. All aerobic organisms including human beings have to counteract to auto-oxidation or

peroxidation using antioxidant defence systems. The antioxidant molecules are obtained from the natural sources as part of food and some of them are also synthesised in the organism itself as part of natural defence against free radicals (3-5). The plants are considered as one of the richest sources of antioxidants and many plants have been reported for their antioxidant properties that contribute for their medicinal value.

*Cardiospermum halicacabum* is a climber plant widely distributed in tropical and sub tropical regions of Asia and Africa. *C. halicacabum* is also found in farmland of Western ghat region as weed and used for various ailments by the traditional healers (6-8). In the present study the antioxidant potential of *Cardiospermum halicacabum* Linn Leaf was evaluated in the atherodiet induced

*\*Address for Corresponding Author:*

Dr. N. Thamizh Selvam Ph.D.,  
Assistant Director- Biochemistry  
National Ayurveda Research Institute for Panchakarma  
(CCRAS, Ministry of AYUSH, Govt. of India)  
Cheruthuruthy, Kerala- 679 531 India.  
Email: nthamizhselvam@gmail.com

DOI: <https://doi.org/10.31024/apj.2018.3.2.3>2456-1436/Copyright © 2018, N.S. Memorial Scientific Research and Education Society. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

hyperlipidemic wistar albino rats.

## Materials and methods

### Chemicals and Reagents

The Chemicals and Reagents of make Transasia Bio-medicals India Ltd., Spectrum India Ltd, Merck India Ltd, Nice India Ltd, were used for the present experiment.

### Instruments

Fully Auto Biochemistry Analyzer EM200- Make-Transasia and UV-Visible Spectrophotometer- Make- Agilent were the major equipments used for the present study.

### Plant Authentication

The plant *Cardiospermum halicacabum* was collected from the Western ghat Region (Palghat and Thrissur). The authentication of the plant was done by Taxonomist, Kerala Forest Research Institute (KFRI), Government of Kerala, Thrissur. Voucher specimen is maintained in the Biochemistry Department of NARIP, Cheruthuruthy.

### Extract Preparation

The leaves of *Cardiospermum halicacabum* were collected and used for the study. The Hydro alcoholic extract of leaf (one part alcohol and one part water) was prepared as per Ayurvedic Pharmacopoeia of India Part I Vol. VIII.

### Ethical Committee approval

Institutional Animal Ethics Committee's (IAEC) approval was obtained for the animal experiments vide Proposal No. IAEC/NRIP/2015-16/05 dated 23.01.2016 in the meeting held at National Research Institute for Panchakarma, Cheruthuruthy, Thrissur, Kerala.

### Acclimatization of animals

Male and Female Wistar Albino Rats in a test group were housed individually in standard polypropylene cages with top grill having facilities for pelleted feed and unlimited supply of drinking water in glass bottles with sipper tubes. An ideal temperature and relative humidity were maintained with a 12hr light / dark cycle. The animals had free access to sterile pelleted feed of standard composition and purified water.

### Preparation of Atherodiet

High fat diet or Atherodiet was prepared as per the method of Bopanna *et al.*, (1997) comprising of 2% Cholesterol, 0.25% Bile salts and 15% Butter were used for the present study.

### Experimental Study Design

A total of 36 Wistar Albino Rats of 12 weeks old were randomized and equally divided into the 6 groups. Each group consisted of 6 animals (3 Males and 3 Females) (Pillai *et al.*, 2012, ThamizhSelvam *et al.*, 2015). Control group received

standard diet and distilled water, Test extract group received 250mg/kgbw *C. halicacabum* extract and standard diet, Disease control group received only Atherodiet, Treatment group-Low dose received Atherodiet with 250mg/kgbw *C. halicacabum* extract, Treatment group-High dose received Atherodiet with 500mg/kgbw *C. halicacabum* extract and Standard drug group received Atherodiet with 5mg/kgbw Atorvastatin (Patil *et al.*, 2010, Adeneye *et al.*, 2009, Hamden *et al.*, 2009, Mariyappan *et al.*, 2011, Maruthappan *et al.*, 2010). The details have been briefed in Table 1.

**Table 1.** Details of Experimental animal groups and their test extract dosages

S. No	Groups	Name of groups	Details
1	Group 1	Control Group	Standard Diet
2	Group 2	Test Extract Group	Standard Diet + <i>C. halicacabum</i> extract 250mg/kgbw
3	Group 3	Disease Control Group	Atherodiet (High fat diet)
4	Group 4	Treatment Group- Low dose	Atherodiet + <i>C. halicacabum</i> extract 250mg/kgbw
5	Group 5	Treatment Group- high dose	Atherodiet + <i>C. halicacabum</i> extract 500mg/kgbw
6	Group 6	Standard Drug Group	Atherodiet + Atorvastatin 5mg/kgbw

### Sample collection and Biochemical Studies

After 4 weeks treatment, body weights were measured and blood samples were collected by retro orbital route after overnight fasting of the animals for the study of biochemical parameters. Blood samples were collected for evaluation of antioxidant enzymes. At the end of the experiment, animals were euthanized and the organs were collected for the experimental purpose.

### Tissue Homogenization

Tissue Homogenization is a process used to prepare tissue samples. 1g tissue was weighed and placed in the homogenizing cup. About 5ml phosphate buffer of pH 7.0 was added to the homogenizing cup and homogenized. Another 5ml quantity of same buffer was added and then homogenized. From this, 2ml of homogenate was transferred for the determination of Glutathione and the remaining part of the homogenate was centrifuged at 10,000 rpm for 20min at 4°C. The supernatant was collected and used for the other enzyme assays after determination of protein concentration.

### Protein Estimation in Tissue

Protein in the tissue and serum/blood were assayed using the Bradford's test. The assay is based on the observation that the absorbance maximum for acidic solution of

Coomassie Brilliant Blue G-250 shifts from 465 to 595 nm when binding to protein occurs. Accordingly the colour change was read at 595nm.

### Catalase

Catalase was estimated by the method of Beers *et al.*, (1952). 0.01ml of the tissue supernatant was mixed with 1.9ml of Phosphate buffer (0.5 M, pH-7.0) and 1ml of 11mM H<sub>2</sub>O<sub>2</sub>. Decrease in absorbance per minute was measured at 240nm for the first three minute against a blank which contained 0.01ml of the tissue supernatant and 2.9ml of Phosphate buffer.

### Glutathione Peroxidase (GPx) activity

Glutathione peroxidase activity was determined according to the method of Hafemann *et al.*, 1974. Tissue homogenate (approximately 0.5 mg protein) was incubated with 0.1 ml of 5mM GSH, 0.1 ml of 1.25 mM H<sub>2</sub>O<sub>2</sub>, 0.1 ml of 25mM NaN<sub>3</sub> and phosphate buffer (0.05mM, pH7) in a total volume of 2.5 ml at 37°C for 10 Min. The absorbance of yellow coloured complex was measured at 412 nm after incubation for 10 min at 37 °C against distilled water. A sample without the tissue homogenate was kept as control.

### Glutathione (GSH)

The GSH was measured by the method of Moron *et al.*, 1979. 0.5 ml of the tissue homogenate was mixed with 0.1 ml of 25 % TCA and kept on ice for few minutes. This was then centrifuged at 2500 rpm for 5 minutes to settle down the precipitate. 0.3 ml of

the supernatant was mixed with 0.7 ml of 0.2M sodium phosphate buffer (pH 8.0) and 2 ml of 0.6 mM DTNB (prepared in 0.2 M phosphate buffer, pH 8.0. The yellow colour obtained was read after 10 minutes at 412 nm against blank.

### Superoxide dismutase (SOD)

Superoxide dismutase activity was determined according to the method of McCord and Fridovich *et al.*, 1969. 0.05 ml of the homogenate was mixed with 0.2 ml of 0.1 M EDTA (containing 0.0015% NaCN), 0.1 ml of 1.5mM NBT and phosphate buffer (67 mM, pH 7.8) in a total volume of 2.6 ml. Percent of inhibition was calculated after comparing absorbance of sample with the absorbance of the control.

### Statistical Analysis

One way ANOVA with Post-hoc analysis was carried out. Dunnett's test was applied to compare the dose groups with control arm.

### Results and discussion


#### Antioxidant parameters in blood

The antioxidant parameters such as Catalase, Glutathione peroxidase, Glutathione and Superoxide dismutase (SOD) were evaluated in blood samples of experimental animals. It is observed that there were enhanced levels of antioxidant parameters in Treatment group Low dose and it was not significant as compared with Disease Control and Healthy


**Table 2.** Antioxidant enzyme parameters in Blood of Wistar albino Rats during Efficacy study.

S. No.	Groups	Antioxidant Enzymes in Blood			
		Catalase (U/mg ptn)	Glutathione peroxidase (U/gHb)	Glutathione (nmol/mg ptn)	Superoxide dismutase (U/gHb)
1	Group 1: Healthy Control Group	44.26 ± 3.71	1135.0 ± 97.70	513.87 ± 27.72	869.62 ± 98.90
2	Group 2: Test Extract Group	43.39 ± 6.05 <sup>a</sup>	1170.41 ± 108.58 <sup>a</sup>	549.91 ± 26.42 <sup>a</sup>	895.96 ± 98.51 <sup>a</sup>
3	Group 3: Disease Control Group	39.32 ± 2.27 <sup>a</sup>	968.57 ± 62.05 <sup>a</sup>	459.91 ± 11.87 <sup>a</sup>	801.79 ± 114.81 <sup>a</sup>
4	Group 4: Treatment Group-Low Dose 250 mg/kgbw	47.91 ± 3.49 <sup>ab</sup>	1215.05 ± 95.35 <sup>ab</sup>	590.62 ± 29.44 <sup>ab*</sup>	995.27 ± 100.78 <sup>ab</sup>
5	Group 5: Treatment Group- High Dose 500 mg/kgbw	76.82 ± 4.27 <sup>a**b**</sup>	1581.86 ± 120.33 <sup>a**b**</sup>	683.37 ± 43.06 <sup>a**b**</sup>	1223.99 ± 101.81 <sup>ab</sup>
6	Group 6: Standard Drug Atorvastatin 5mg/kgbw	46.06 ± 3.40 <sup>ab</sup>	1222.70 ± 61.54 <sup>ab</sup>	491.33 ± 23.34 <sup>ab</sup>	730.74 ± 109.31 <sup>ab</sup>

Values are expressed as Mean ± SEM, N=6 per group. Values with superscript a, b indicate no significant difference (p>0.05) when compared with Healthy Control, Disease Control respectively. Values with superscript a\*/a\*\* indicate significant difference compared with Healthy Control; b\*/b\*\* indicate significant difference compared with Disease Control at p<0.05 and p<0.01 respectively



**Figure 1.** Effect of *C. halicacabum* extract on Catalase and GSH in Blood




**Figure 2.** Effect of *C. halicacabum* extract on GPx and SOD in Blood

Control. The Treatment Group High dose showed that there were significant enhancement of Catalase, Glutathione peroxidase and Glutathione levels as compared with Disease Control and Healthy Control ( $p < 0.01$ ). Superoxide dismutase level is found to be increased in High dose group but was not significant.


Atorvastatin treated Standard Drug group also showed elevated levels of antioxidant enzymes as compared to Healthy Control and Disease Control but it was not significant (Table 2 and Figure 1-2).

**Antioxidant parameters in tissues**

The antioxidant parameters in tissue sample of Liver, Kidney and Heart were analyzed. The antioxidant enzymes Catalase, Glutathione peroxidase, Glutathione and



**Figure 3.** Effect of *C. halicacabum* extract on Catalase and GPx in Liver



**Figure 4.** Effect of *C. halicacabum* extract on GSH and SOD in Liver

**Table 3.** Antioxidant enzyme parameters in liver tissue of wistar albino rats during Efficacy study


S. No.	Groups	Antioxidant Enzymes in Tissues - Liver			
		Catalase (U/mg ptn)	Glutathione peroxidase (U/mg ptn)	Glutathione (nmol/mg ptn)	Superoxide dismutase (U/mg ptn)
1	Group 1: Healthy Control Group	143.47 ± 4.46	77.78 ± 8.47	4.74 ± 0.30	4.86 ± 0.17
2	Group 2: Test Extract Group	148.41 ± 6.50 <sup>a</sup>	80.63 ± 3.57 <sup>a</sup>	5.41 ± 0.26 <sup>a</sup>	5.43 ± 0.31 <sup>a</sup>
3	Group 3: Disease Control Group	123.25 ± 3.96 <sup>a</sup>	66.09 ± 3.98 <sup>a</sup>	3.98 ± 0.07 <sup>a</sup>	4.00 ± 0.22 <sup>a</sup>
4	Group 4: Treatment Group-Low Dose 250 mg/kgbw	161.69 ± 6.85 <sup>ab**</sup>	85.75 ± 3.25 <sup>ab**</sup>	5.53 ± 0.27 <sup>ab**</sup>	5.40 ± 0.35 <sup>ab**</sup>
5	Group 5: Treatment Group- High Dose 500 mg/kgbw	193.58 ± 6.59 <sup>a**b**</sup>	95.18 ± 3.63 <sup>ab**</sup>	6.71 ± 0.12 <sup>a**b**</sup>	6.80 ± 0.12 <sup>a**b**</sup>
6	Group 6: Standard Drug Atorvastatin 5mg/kgbw	154.72 ± 3.93 <sup>ab**</sup>	77.99 ± 3.19 <sup>ab</sup>	5.06 ± 0.20 <sup>ab**</sup>	4.29 ± 0.31 <sup>ab</sup>

Values are expressed as Mean ± SEM, N=6 per group. Values with superscript a, b indicate no significant difference ( $p > 0.05$ ) when compared with Healthy Control, Disease Control respectively. Values with superscript a\*/a\*\* indicate significant difference compared with Healthy Control; b\*/b\*\* indicate significant difference compared with Disease Control at  $p < 0.05$  and  $p < 0.01$  respectively


**Table 4.** Antioxidant enzyme parameters in Kidney Tissue of Wistar Albino Rats during Efficacy study

S. No.	Groups	Antioxidant Enzymes in Tissues - Kidney			
		Catalase (U/mg ptn)	Glutathione peroxidase (U/mg ptn)	Glutathione (nmol/mg ptn)	Superoxide dismutase (U/mg ptn)
1	Group 1: Healthy Control Group	125.18 ± 7.82	89.02 ± 1.42	17.01 ± 0.47	5.01 ± 0.19
2	Group 2: Test Extract Group	129.01 ± 3.72 <sup>a</sup>	105.82 ± 3.37 <sup>a*</sup>	18.23 ± 0.81 <sup>a</sup>	5.54 ± 0.22 <sup>a</sup>
3	Group 3: Disease Control Group	115.80 ± 2.81 <sup>a</sup>	80.81 ± 3.06 <sup>a</sup>	13.54 ± 0.55 <sup>a**</sup>	3.70 ± 0.21 <sup>a*</sup>
4	Group 4: treatment Group-Low Dose 250 mg/kgbw	130.79 ± 8.16 <sup>ab</sup>	108.09 ± 6.36 <sup>a*b**</sup>	18.35 ± 0.69 <sup>ab**</sup>	5.98 ± 0.14 <sup>ab**</sup>
5	Group 5: treatment Group- High Dose 500 mg/kgbw	145.41 ± 5.53 <sup>ab**</sup>	127.43 ± 2.86 <sup>a**b**</sup>	22.68 ± 1.10 <sup>a**b**</sup>	7.03 ± 0.32 <sup>a**b**</sup>
6	Group 6: Standard Drug Atorvastatin 5mg/kgbw	110.36 ± 2.26 <sup>ab</sup>	100.34 ± 5.78 <sup>ab*</sup>	13.65 ± 0.52 <sup>a*b</sup>	4.62 ± 0.58 <sup>ab</sup>

Values are expressed as Mean ± SEM, N=6 per group. Values with superscript a, b indicate no significant difference (p>0.05) when compared with Healthy Control, Disease Control respectively. Values with superscript a\*/a\*\* indicate significant difference compared with Healthy Control; b\*/b\*\* indicate significant difference compared with Disease Control at p<0.05 and p<0.01 respectively




**Figure 5.** Effect of *C. halicacabum* extract on Catalase and GPx in Kidney




**Figure 6.** Effect of *C. halicacabum* extract on GSH and SOD in Kidney

Superoxide dismutase levels in liver tissues of Extract treated rats showed significant increase when compared with Disease Control (p<0.01). The Atorvastatin treated Standard Drug group also showed significant improvement in antioxidant profile when compared to Disease Control (Table 3 & Figure 3-4). The antioxidant enzymes Catalase and Glutathione peroxidase levels in kidney tissue samples of Disease Control group showed no significant changes when compared with Healthy Control. But, the decreased levels of Glutathione and Superoxide dismutase

levels were noted when compared with Healthy Control (p<0.01 and p<0.05) respectively. The kidney tissues of Treatment group Low dose showed there was significant enhancement (p<0.01) in Glutathione peroxidase, Glutathione and Superoxide dismutase levels but no significant changes observed in Catalase as compared to Healthy Control and Disease Control. The Treatment group High Dose showed elevated levels of all antioxidant parameters as compared to Disease Control (p<0.01) and



**Figure 7.** Effect of *C. halicacabum* extract on Cat.andGPx in Hea



**Figure 8.** Effect of *C. halicacabum* extract on GSH and SOD in Heart

**Table 5.** Antioxidant enzyme parameters in Heart Tissue of Wistar Albino Rats during Efficacy study

S. No.	Groups	Antioxidant Enzymes in Tissues - Heart				
		Catalase (U/mg ptn)	Glutathione (U/mg ptn)	peroxidase (U/mg ptn)	Glutathione (nmol/mg ptn)	Superoxide dismutase (U/mg ptn)
1	Group 1: Healthy Control Group	13.30 ± 1.06	92.48 ± 2.32		3.04 ± 0.12	6.42 ± 0.23
2	Group 2: Test Extract Group	14.05 ± 0.83 <sup>a</sup>	96.60 ± 2.35 <sup>a</sup>		3.23 ± 0.14 <sup>a</sup>	6.49 ± 0.24 <sup>a</sup>
3	Group 3: Disease Control Group	10.07 ± 0.62 <sup>a*</sup>	80.37 ± 4.06 <sup>a</sup>		2.27 ± 0.03 <sup>a*</sup>	5.77 ± 0.25 <sup>a</sup>
4	Group 4: reatment Group- Low Dose 250 mg/kgbw	14.14 ± 0.67 <sup>ab**</sup>	98.22 ± 4.56 <sup>ab*</sup>		3.20 ± 0.16 <sup>ab*</sup>	7.23 ± 0.18 <sup>ab**</sup>
5	Group 5: reatment Group- High Dose 500 mg/kgbw	17.22 ± 0.75 <sup>a**b**</sup>	106.64 ± 2.88 <sup>a**b**</sup>		3.61 ± 0.14 <sup>ab**</sup>	8.52 ± 0.40 <sup>a**b**</sup>
6	Group 6: Standard Drug Atorvastatin 5mg/kgbw	13.19 ± 0.79 <sup>ab*</sup>	83.20 ± 3.17 <sup>ab</sup>		3.12 ± 0.38 <sup>ab</sup>	7.06 ± 0.25 <sup>ab*</sup>

Values are expressed as Mean ± SEM, N=6 per group. Values with superscript a, b indicate no significant difference (p>0.05) when compared with Healthy Control, Disease Control respectively. Values with superscript a\*/a\*\* indicate significant difference compared with Healthy Control; b\*/b\*\* indicate significant difference compared with Disease Control at p<0.05 and p<0.01 respectively.

Healthy Control (p<0.01). The Atorvastatin treated group showed significant variations in Glutathione peroxidase and Glutathione levels and no significant changes in Catalase and Superoxide dismutase levels as compared to Healthy Control and Disease Control (Table 4 & Figure 5-6). The antioxidant parameters such as Catalase, Glutathione peroxidase, Glutathione and Superoxide dismutase levels in Heart tissue samples of treatment groups i.e. Low dose and High dose group showed enhanced levels as compared with Disease Control group. The Atorvastatin treated group showed significant level (p<0.05) of increase in Catalase and Superoxide dismutase level when compared with Disease Control group and no significant changes in Glutathione peroxidase and Glutathione levels. (Table 5 and Figure 7-8).

### Conclusion

The study showed that the Hydro alcoholic extract of *C. halicacabum* Linn Leaf has significant hypolipidemic activity in the atherodiet induced hyperlipidemia in Wistar rats. It also demonstrated that the antioxidant parameters such as Catalase, Glutathione peroxidase, Glutathione and Superoxide dismutase levels were found significantly increased in the blood and tissue samples. So, the study concluded that *C. halicacabum* extract has significant antioxidant potential and that must be contributing for its medicinal value.

### Acknowledgement

Authors are thankful to the Director General, CCRAS, New Delhi for the Intra Mural Research Project Grant. The authors also expressing the gratitude to Laboratory Technicians of Biochemistry and Pathology Department Mr. Venugopalan TN,

Mr.Sanal Gopi CG, Mrs..Ranjini KR and Mrs. Neethu KR for their extended help in the study.

**Conflict of interest:** Nil

### References

- Adeneye AA, Olagunju. 2009. Preliminary Hypoglycemic and Hypolipidemic activities of the aqueous seed extract of *Carica papaya* Linn in Wistar rats. *Biology and Medicine*, 1(1):1-10.
- Beers RF, Sizer IW. 1952. Spectrophotometric method for measuring the breakdown of Hydrogen Peroxide by Catalase. *Journal of Biological Chemistry*, 195:133-140.
- Bopanna KN, Bhagyalakshmi N, Rathod SP, Balaraman R, Kannan J. 1997. Cell culture derived *Hemidesmus indicus* in the prevention of Hypercholesterolemia in Normal and Hyperlipidemic rats. *Indian journal of Pharmacology*, 29:105-109.
- Glenson HA. 1991. *Manual of Vascular Plants of Northeastern United States and Adjacent Canada*. New York, USA: The New York Botanical Garden.
- Hafeman DG, Sunde RA, Hoekstra WG. 1974. Effect of Dietary Selenium on Erythrocyte and Liver Glutathione peroxidase in the Rat. *Journal of Nutrition*, 104(5):580-587.
- Hamden K, Allouche N, Damak M, Elfeki A. 2009. Hypoglycemic and antioxidant effects of phenolic extracts and purified hydroxyl tyrosol from olive mill waste in vitro in rats. *Chemico-Biological Interactions*,

- 180:421-432.
- Harman D. 1992. Role of free radicals in ageing and disease. *Annals of the New York Academy of Sciences*, 673:598-620.
- Johnston SK, Murray DS, Williams JC. 1979. Germination and emergence of balloon vine (*Cardiospermum halicacabum*). *Weed Science*, 27(1):73-76.
- Mariyappan M, Bharathidasan R, Madhanraj P, Paneerselvam A, Ambikapathy V. 2011. Antibacterial activity of *Cardiospermum halicacabum* and *Melothria heterophylla*. *Asian Journal of Pharmaceutical Research*, 1(4):111-113.
- Maruthappan V, Sakthi Shree K. 2010. Hypolipidemic activity of Haritaki (*Terminalia chebula*) in Atherogenic Diet induced Hyperlipidemic Rats. *Journal of Advanced Pharmaceutical Technology & Research*, 1(2):229-235.
- McCord JM, Fridovich I. 1969. Superoxide Dismutase, An enzymic function for Erythrocyte (Hemocuprein). *The Journal of Biological Chemistry*, 244(22):6049-6055.
- Moron MS, Depierre JW, Mannervik B. 1979. Levels of Glutathione, Glutathione reductase and Glutathione S-transferase activities in rat lung and liver. *Biochem Biophys Acta*, 582(1):67-78.
- Patil RH, Prakash K, Maheswari VL. 2010. Hypolipidemic effect of *Celastrus paniculatus* in experimentally induced Hypercholesterolemic Wistar rats. *Indian Journal of Clinical Biochemistry*, 25(4):405-410.
- Pawan K, Dhirender K, Sukhbir LK. 2011. In vivo antioxidant activity of plant *Abutilon indicum*. *Journal of Pharmaceutical Education Research*, 2(1):50-53.
- Pillai NR, Vijayamma N. 1985. Some Pharmacological studies on *Cardiospermum alicacabum* Linn. *Ancient Science of Life*, 1(7):32-36.
- Prior RL, Cao G. 2000. Analysis of botanicals and dietary supplements for antioxidant capacity: A review. *The Journal of AOAC International*, 83:950-956.
- Romero FJ, Morell FB, Romero MJ, Jareno EJ, Romero B, Marin N. 1998. Lipid peroxidation products and antioxidants in human disease. *Environmental Health Perspective*, 106:1229-1234.
- ThamizhSelvam N, Prasannakumari K, Sanjayakumar YR, Surabi KR, Venugopalan TN, Vasanthakumar KG, Acharya MV. 2015. Evaluation of Hypocholesterimic Activity of SPHAG – A poly Herbal formulation in Wistar Albino Rats. *International Journal for Pharma Sciences and Research*, 6(10):1245-1249.
- Warrier PK, Nambiar VPK, Ramankutty C. 2002. *Indian Medicinal Plants: A compendium of 500 species*. Orient Longman Publications, Vol1:377-379.
- Yamaguchi F, Saito M, Ariga T, Yoshimura Y, Nakazawa H. 2000. Free radical scavenging activity of garcinol from *Garcinia* fruit rind. *Journal of Agricultural and Food Chemistry*, 48:2320-2325.